

Vaccination and epilepsy

In this leaflet Dr John Paul Leach, a consultant neurologist in Glasgow, talks about vaccinations for people with epilepsy.



Dr John Paul Leach

I have epilepsy

Does this change what vaccinations I can and can't receive?

People who have epilepsy can safely be vaccinated with almost all vaccines. Epilepsy itself is not a contraindication for vaccination. If you regularly have seizures, or are taking any type(s) of anti-epileptic drug, you should still be vaccinated absolutely as normal.

Your children should also be vaccinated as normal. A family history of epilepsy does not affect vaccination in any way.

> You should not receive a vaccine if you have had a clear and confirmed allergic reaction to the same vaccine or another vaccine against the same disease.

If you have an allergy to eggs you should not receive influenza or yellow fever vaccines, as these contain egg proteins.

If you are allergic to eggs and require these vaccines, your doctor will seek specialist advice about vaccinating you under expert supervision.

My child has epilepsy

Should they receive the normal childhood vaccines?

Children with epilepsy can safely be vaccinated if their condition is stable. Epilepsy itself is not a contraindication for vaccination.

Very young children (under 12 months old) who have epilepsy may have their vaccination postponed while their condition is investigated and stabilised.

Continued overleaf...



...continued from overleaf

Some vaccines can cause a short rise in body temperature as a side effect. Rarely, this can trigger a seizure (called a febrile seizure). If your child has had a febrile seizure in the past, or if anyone in your family has a history of febrile seizures, then they have a slightly increased risk of having a seizure after their vaccination. However they should still be vaccinated as normal.

Ask for advice about how to prevent and manage a high body temperature in your child when they receive the vaccine. For example, giving regular paracetamol for 24 hours after vaccination may be advisable for some children receiving the DTP vaccine.

Vaccines that may cause a temperature include the DTP vaccine, and the measles, mumps and rubella (MMR) vaccine. The high temperature usually happens within 24 hours with the DTP vaccine, but may take up to a week to develop with the MMR vaccine.

.....

This information is taken from:

• "The Green Book", UK Department of Health information on Immunisation Against Infectious Disease. 2006 edition.

The latest version is available to download from: www.dh.gov.uk/greenbook

• BMJ Best Health website (<http://besthealth.bmj.com/x/index.html>)

This is one in a series of information leaflets about epilepsy.

To access the others, or to find out more about our research, please visit our website: www.epilepsyresearch.org.uk

Text adapted from material supplied by Epilepsy Connections, February 2014.



**epilepsy
research uk**

Epilepsy Research UK
PO Box 3004 London W4 4XT

- ☎ 020 8747 5024
- ☎ 0870 838 1069
- ✉ info@eruk.org.uk
- 🌐 www.epilepsyresearch.org.uk

Registered office: Lamb House, Church Street, London W4 2PD. Registered charity no: 1100394

